



Patient Sticker

<p>Date of Epidural placement:</p> <p>Epidural level:</p> <p>Epidural Medication:</p> <p><input type="checkbox"/> Bupivacaine 0.1%</p> <p><input type="checkbox"/> Fentanyl 2mcg/mL with Bupivacaine 0.0625%</p> <p><input type="checkbox"/> Morphine 50mcg/mL with Bupivacaine 0.0625%</p> <p><input type="checkbox"/> Hydromorphone 20mcg/mL with Bupivacaine 0.0625%</p> <p>Other: _____</p> <p>Rate: _____ mL/hr</p> <p>Boluses: ____ mL every ____ minutes for a total of ____ times per hour</p> <p>Foley necessary for epidural: Yes No</p>	<p>MD/RN communication/patient concerns:</p> <ul style="list-style-type: none"> Do not let epidural bag or tubing run dry Patient's abdomen and legs may feel numb from epidural medication infusion. Page 4878 for the following signs and symptoms: <ul style="list-style-type: none"> - SBP less than ____ mmHg or HR less than ____. - Continuous leaking past POD#1. - Increased loss of motor function in lower extremities, or decreased sensation or weakness in upper extremity <p style="color: red; font-weight: bold; margin-top: 10px;"> Do not change or discontinue epidural pump settings or orders unless discussed with Acute Pain Team first. Notify Acute Pain Team if epidural paused, disconnected or epidural order set missing from MAR. </p>
--	--

Questions? Please page Acute Pain Pager 4878.
Emergency? Call 50250 or 51354.

NOT PART OF THE MEDICAL RECORD